

## Texas Dairy Matters

Higher Education Supporting the Industry

## **Prussic Acid in Forages**

Tamilee Nennich, Ph.D.
Extension Dairy Specialist
Department of Animal Science
Texas A&M AgriLife xtension Service
The Texas A&M University System

Under certain conditions prussic acid (HCN or hydrocyanic acid) accumulates in sorghum, sudangrass, johnsongrass, and sorghum-sudangrass hybrids. Feeding forages containing prussic acid to livestock can be fatal.

After forages containing prussic acid are consumed, the cyanide ion binds to the oxygen in the blood and prevents oxygen from being released from the hemoglobin to body tissues. The bound oxygen in the blood results in bright cherry-red colored blood that is characteristic of prussic acid poisoning.

Prussic acid is produced in plants under stress conditions. High rates of N fertilization can increase prussic acid accumulation in forages. Typically, the leaves accumulate more prussic acid than do stems. The greater leaf-to-stem ratios that occur during drought conditions increase the chance of prussic acid poisoning. In addition, re-growth after a rain is often associated with prussic acid poisoning.

Test forages to determine the levels of prussic acid. Prussic acid levels less than 500 ppm of CN as HCN on a dry basis are considered safe. When prussic acid levels are greater than 1000 ppm on a dry basis, the forage is considered hazardous. Do not feed forage with high levels of prussic acid

When prussic acid levels are a concern in forages, consider either ensiling or haying the forage. The curing and ensiling processes decrease the prussic acid concentrations and make the forage safe to feed to dairy cattle.

http://texasdairymatters.org

July, 2005

The Texas A&M AgriLife Extension Service provides equal opportunities in its programs and employment to all persons, regardless of race, color, sex, religion, national origin, disability, age, genetic information, veteran status, sexual orientation, or gender identity.