

Texas Dairy Matters

Higher Education Supporting the Industry

In this updated article, the authors corrected an error where the 2.36 mm BPS was mistakenly labeled as 1.7 mm BPS. The authors regret the mistake and apologize for the oversight.

Strategies to include sorghum silage in lactating cow rations. Part 3.

Juan M. Piñeiro, DVM, MS, Ph.D., Douglas Duhatschek, DVM, Artur Grando Pilati, DVM
 Department of Animal Science, The Texas A&M University System
Guest authors: Lyndon Luckasson, OEM Business Director, Scherer Inc.,
 Elizabeth Coons, Research and Development Coordinator, Rock River Laboratory

Introduction

The expansion of dairy cattle inventory in the Texas Panhandle is increasing forage demand while groundwater levels decline. Sorghum is a drought-tolerant forage alternative with low input costs, and production of sorghum for silage has increased in recent years in the U.S. and Texas (Fig. 1)¹. However, conventional sorghum silage has lower fiber and starch digestibility compared to corn. In a previous [article](#), we shared preliminary results of a new grain processing technology that significantly increases berry processing². This article will share the berry processing score and starch digestibility of sorghum silage from the same experiment.




Figure 1. Production of sorghum for silage in the U.S. and Texas. From 2020 to 2023, production of sorghum silage increased by 59% in the U.S. and 92% in Texas. Data from USDA-NASS. 2004-2024.

A new processing technology may be a game changer for sorghum silage

In collaboration with Scherer Inc., we randomly collected duplicated samples from three different areas of a field with a forage sorghum hybrid (Pearl, MOJO seeds, Hereford, Texas) under center pivot irrigation harvested at the hard dough stage³. The objectives were to assess the effect of a novel kernel processing, KP, technology and ensiling time on sorghum silage processing and rumen in situ starch digestibility after seven hours of incubation. The experiment consisted of two different ensiling times -- 0 or 90 days -- and three different harvesting strategies:

- 1- Low noKP: Low cut height (8 in) without KP.
- 2- Low +KP: Low cut height (8 in) with KP.
- 3- High +KP: High cut height (48 in) with KP.

Compared to Low noKP, using KP significantly increases the berry processing score, BPS, (percentage of starch passing a 2.36 mm sieve, Table 1)³. Noticeably, Low +KP achieved a BPS two times higher than the 28% suggested as an initial goal by other authors⁴.

Sorghum silage harvested with this outstanding berry processing combined with a fermentation length of 90 days achieved a remarkable rumen in-situ starch digestibility. Regardless of cut height, using KP and fermentation length of 90 days roughly doubled the in-situ starch digestibility compared to not using KP. Noticeably, the in-situ starch digestibility after seven hours of incubation increased after ensiling both cut heights when harvesting with KP and ensiling for 90 days but not without KP. This may happen due to the increased access of bacteria to starch during the ensiling time in the treatments with KP where berry processing was successful whereas the pericarp would impede bacterial access to the starch in the Low noKP treatment.

| Item | Low noKP | | Low +KP | | High +KP | | SEM | P-value | | |
|-----------------------|-------------------|-------------------|--------------------|--------------------|-------------------|--------------------|------|---------|-------|-------|
| | 0 d | 90 d | 0 d | 90 d | 0 d | 90 d | | HS | ET | HS*ET |
| ADF, % DM | 24.3 ^a | 23.9 ^a | 22.1 ^a | 22.0 ^{ab} | 15.9 ^c | 16.8 ^{bc} | 1.13 | <0.01 | 0.91 | 0.85 |
| aNDFom, % DM | 34.2 ^a | 33.3 ^a | 30.4 ^{ab} | 27.5 ^b | 22.2 ^c | 22.4 ^c | 1.05 | <0.01 | 0.14 | 0.25 |
| Starch, % DM | 33.1 ^b | 30.6 ^b | 37.2 ^b | 36.7 ^b | 52.1 ^a | 51.2 ^a | 1.67 | <0.01 | 0.29 | 0.76 |
| 2.36 mm BPS, % starch | 7.7 ^c | 8.6 ^c | 55.8 ^b | 56.7 ^b | 70.1 ^a | 71.6 ^a | 1.28 | <0.01 | 0.22 | 0.93 |
| isSD7, % starch | 29.0 ^b | 25.8 ^b | 39.9 ^b | 65.9 ^a | 36.2 ^b | 66.7 ^a | 4.62 | <0.01 | <0.01 | <0.01 |

Table 1. Nutrient composition, berry processing score (BPS), and rumen in situ starch digestibility after 7 hours of incubation (isSD7) of sorghum silage with two different ensiling times (0 or 90 days) and processed with three different harvesting strategies (Low noKP: low cut height without KP, Low +KP: low cut height with KP, High +KP: high cut height with KP). Numbers with different superscripts are significantly different.

Increasing the cut height did not increase in situ starch digestibility

We hypothesized that increasing the cut height from 8 inches to 48 inches, to increase the sorghum panicle to stover proportion, would increase berry processing and in situ starch digestibility. Compared to Low +KP, High +KP increased the BPS by roughly 26%. However, there was no significant difference between Low +KP and High +KP on in situ starch digestibility (Fig. 2). Perhaps this could be attributed to both treatments achieving a very good BPS. In this study, the BPS of Low +KP and High +KP were roughly 56% and 71%.




Figure 2. The effect of increasing the cut height from 0.2 to 1.2 m to increase the panicle to vegetative plant parts ratio on berry processing score (BPS) and rumen in-situ starch digestibility (isSD7). Compared to harvesting with low cut height without KP (Low +KP), harvesting with a high cut height with KP (High +KP) did not increase isSD7.

Our results suggest this new processing technology combined with a 90-day ensiling time increases sorghum berry processing and in-situ starch digestibility, and increasing cut height may increase BPS but may not further increase in-situ starch digestibility. The remarkable 66% of rumen in-situ starch digestibility after seven hours of incubation achieved is 12% below the 75% average from Rock River Lab for corn silage in the U.S⁵. Future studies should assess the effect of further increasing ensiling time in sorghum silage harvested with this technology on starch digestibility.

Acknowledgements

We would like to acknowledge Sherer Inc. and Rock River Laboratory for their support and collaboration with this ongoing research as well as dairy farmers and students who participated in the field data collection and curation.

References

- ¹ USDA-NASS. 2004-2024. Crop production 2003 to 2023 Summary reports.
- ² Piñeiro J.M., D. Duhatschek, A.G. Pilati, L. Luckasson. 2024. Strategies to include sorghum silage in lactating cows rations. Part 2. Texas Dairy Matters.
- ³ D. Duhatschek, A. Grando Pilati, L. Luckasson, J. Goeser, E. Coons, L.F. Ferraretto, J.M. Piñeiro. 2024. The effects of a new grain processing technology on forage sorghum processing. J. Dairy Sci. (accepted for publication).
- ⁴ Raver K., J. Goeser, S. Marshall. 2023. A survey of berry processing score and nutrient content of sorghum silage on commercial livestock operations across the US. J. Anim. Sci.
- ⁵ Rock River Laboratory, 2023. Corn Silage isSD7 data from samples submitted to Rock River Laboratory between September 29th, 2023 and November 30, 2023.

<http://texasdairymatters.org>

May 2024

The Texas A&M AgriLife Extension Service provides equal opportunities in its programs and employment to all persons, regardless of race, color, sex, religion, national origin, disability, age, genetic information, veteran status, sexual orientation, or gender identity.

The Texas A&M University System, U.S. Department of Agriculture, and the County Commissioners Courts of Texas